

Generic Structural Stability in 2×2 Systems of Conservation Laws

Hong Kiat TAN

University of California, Los Angeles

25 Apr 2025


UCLA College | Physical Sciences
Mathematics

Motivation

Gravity-Driven Particle-Laden Flow

Experimental Setup:

- Inclined Slope
- Negatively buoyant monodisperse particles (i.e. glass beads) mixed with oil.
- Mixture of particles and oil added with a gate before the start of the experiment.
- Release the gate to start.



Motivation

$h(x, t)$: Height of the slurry mixture.

$\phi_0(x, t)$: z -averaged particle volume fraction.

x : Distance downstream (from the gate).


Motivation

$h(x, t)$: Height of the slurry mixture.

$\phi_0(x, t)$: z -averaged particle volume fraction.

x : Distance downstream (from the gate).

Assumptions: Fast Equilibrium + Lubrication Assumption.



Motivation

$h(x, t)$: Height of the slurry mixture.

$\phi_0(x, t)$: z -averaged particle volume fraction.

x : Distance downstream (from the gate).

Assumptions: Fast Equilibrium + Lubrication Assumption.

- Conservation of suspension volume:

$$\partial_t h + \partial_x F(h, h\phi_0) = 0.$$


- Conservation of the number of particles:

$$\partial_t (h\phi_0) + \partial_x G(h, h\phi_0) = 0.$$

- Functional form of flux functions:

$$F(h, h\phi_0) = h^3 f\left(\frac{h\phi_0}{h}\right) = h^3 f(\phi_0),$$

$$G(h, h\phi_0) = h^3 g\left(\frac{h\phi_0}{h}\right) = h^3 g(\phi_0).$$



Issue: f and g are computationally expensive to evaluate.

Motivation

To evaluate $f(\phi_0)$ and $g(\phi_0)$ at a single point ϕ_0 , one has to

To evaluate $f(\phi_0)$ and $g(\phi_0)$ at a single point ϕ_0 , one has to

1. Numerically solve the following nonlinear ODE for

$(\phi(s), \sigma(s))$, $s \in [0, 1]$:

$$\begin{cases} \phi'(s) = \frac{(-B_2 + (B_2 + 1)\phi(s) + \rho_s \phi(s)^2)(\phi_m - \phi(s))}{\sigma(s)(\phi_m + (B_1 - 1)\phi(s))} H(\phi(s)) H(\phi_m - \phi(s)), \\ \sigma'(s) = -1 - \rho_s \phi(s), \\ \sigma(0) = 1 + \rho_s \phi_0, \\ \sigma(1) = 0, \end{cases}$$

To evaluate $f(\phi_0)$ and $g(\phi_0)$ at a single point ϕ_0 , one has to

1. Numerically solve the following nonlinear ODE for

$(\phi(s), \sigma(s)), s \in [0, 1]$:

$$\begin{cases} \phi'(s) = \frac{(-B_2 + (B_2 + 1)\phi(s) + \rho_s \phi(s)^2)(\phi_m - \phi(s))}{\sigma(s)(\phi_m + (B_1 - 1)\phi(s))} H(\phi(s)) H(\phi_m - \phi(s)), \\ \sigma'(s) = -1 - \rho_s \phi(s), \\ \sigma(0) = 1 + \rho_s \phi_0, \\ \sigma(1) = 0, \end{cases}$$

2. Compute velocity using $u(s) = \mu_l \int_0^s \sigma(s) \left(1 - \frac{\phi(s)}{\phi_m}\right)^2 ds$.

To evaluate $f(\phi_0)$ and $g(\phi_0)$ at a single point ϕ_0 , one has to

1. Numerically solve the following nonlinear ODE for

$(\phi(s), \sigma(s)), s \in [0, 1]$:

$$\begin{cases} \phi'(s) = \frac{(-B_2 + (B_2 + 1)\phi(s) + \rho_s \phi(s)^2)(\phi_m - \phi(s))}{\sigma(s)(\phi_m + (B_1 - 1)\phi(s))} H(\phi(s)) H(\phi_m - \phi(s)), \\ \sigma'(s) = -1 - \rho_s \phi(s), \\ \sigma(0) = 1 + \rho_s \phi_0, \\ \sigma(1) = 0, \end{cases}$$

2. Compute velocity using $u(s) = \mu_l \int_0^s \sigma(s) \left(1 - \frac{\phi(s)}{\phi_m}\right)^2 ds$.
3. Compute $f(\phi_0) = \int_0^1 u(s) ds$.

To evaluate $f(\phi_0)$ and $g(\phi_0)$ at a single point ϕ_0 , one has to

1. Numerically solve the following nonlinear ODE for

$(\phi(s), \sigma(s)), s \in [0, 1]$:

$$\begin{cases} \phi'(s) = \frac{(-B_2 + (B_2 + 1)\phi(s) + \rho_s \phi(s)^2)(\phi_m - \phi(s))}{\sigma(s)(\phi_m + (B_1 - 1)\phi(s))} H(\phi(s)) H(\phi_m - \phi(s)), \\ \sigma'(s) = -1 - \rho_s \phi(s), \\ \sigma(0) = 1 + \rho_s \phi_0, \\ \sigma(1) = 0, \end{cases}$$

2. Compute velocity using $u(s) = \mu_l \int_0^s \sigma(s) \left(1 - \frac{\phi(s)}{\phi_m}\right)^2 ds$.
3. Compute $f(\phi_0) = \int_0^1 u(s) ds$.
4. Compute $g(\phi_0) = \int_0^1 u(s) \phi(s) ds$.

To evaluate $f(\phi_0)$ and $g(\phi_0)$ at a single point ϕ_0 , one has to

1. Numerically solve the following nonlinear ODE for

$(\phi(s), \sigma(s))$, $s \in [0, 1]$:

$$\begin{cases} \phi'(s) = \frac{(-B_2 + (B_2 + 1)\phi(s) + \rho_s \phi(s)^2)(\phi_m - \phi(s))}{\sigma(s)(\phi_m + (B_1 - 1)\phi(s))} H(\phi(s)) H(\phi_m - \phi(s)), \\ \sigma'(s) = -1 - \rho_s \phi(s), \\ \sigma(0) = 1 + \rho_s \phi_0, \\ \sigma(1) = 0, \end{cases}$$

2. Compute velocity using $u(s) = \mu_l \int_0^s \sigma(s) \left(1 - \frac{\phi(s)}{\phi_m}\right)^2 ds$.
3. Compute $f(\phi_0) = \int_0^1 u(s) ds$.
4. Compute $g(\phi_0) = \int_0^1 u(s) \phi(s) ds$.

Issue: f and g are computationally expensive to evaluate.

Problem Statement

Given:

$$\begin{cases} u_t + (F_0(u, v))_x = 0, \\ v_t + (G_0(u, v))_x = 0, \end{cases}$$

with given initial data
 $(u, v)(0, x)$.

Problem Statement

Given:

$$\begin{cases} u_t + (F_0(u, v))_x = 0, \\ v_t + (G_0(u, v))_x = 0, \end{cases}$$

with given initial data
 $(u, v)(0, x)$.

Can we approximate (F, G)
with (\tilde{F}, \tilde{G}) such that

$$\begin{cases} \tilde{u}_t + (\tilde{F}(\tilde{u}, \tilde{v}))_x = 0, \\ \tilde{v}_t + (\tilde{G}(\tilde{u}, \tilde{v}))_x = 0, \end{cases}$$

with the same initial data
 $(\tilde{u}, \tilde{v})(0, x)$

Problem Statement

Given:

$$\begin{cases} u_t + (F_0(u, v))_x = 0, \\ v_t + (G_0(u, v))_x = 0, \end{cases}$$

with given initial data
 $(u, v)(0, x)$.

Can we approximate (F, G)
with (\tilde{F}, \tilde{G}) such that

$$\begin{cases} \tilde{u}_t + (\tilde{F}(\tilde{u}, \tilde{v}))_x = 0, \\ \tilde{v}_t + (\tilde{G}(\tilde{u}, \tilde{v}))_x = 0, \end{cases}$$

with the same initial data
 $(\tilde{u}, \tilde{v})(0, x)$

yield solutions are sufficiently close in the following sense:

Problem Statement

Given:

$$\begin{cases} u_t + (F_0(u, v))_x = 0, \\ v_t + (G_0(u, v))_x = 0, \end{cases}$$

with given initial data
 $(u, v)(0, x)$.

Can we approximate (F, G)
with (\tilde{F}, \tilde{G}) such that

$$\begin{cases} \tilde{u}_t + (\tilde{F}(\tilde{u}, \tilde{v}))_x = 0, \\ \tilde{v}_t + (\tilde{G}(\tilde{u}, \tilde{v}))_x = 0, \end{cases}$$

with the same initial data
 $(\tilde{u}, \tilde{v})(0, x)$

yield solutions are sufficiently close in the following sense:

- L^1 stability of $L^1 \cap BV$ solution with respect to flux functions, and

Problem Statement

Given:

$$\begin{cases} u_t + (F_0(u, v))_x = 0, \\ v_t + (G_0(u, v))_x = 0, \end{cases}$$

with given initial data
 $(u, v)(0, x)$.

Can we approximate (F, G)
with (\tilde{F}, \tilde{G}) such that

$$\begin{cases} \tilde{u}_t + (\tilde{F}(\tilde{u}, \tilde{v}))_x = 0, \\ \tilde{v}_t + (\tilde{G}(\tilde{u}, \tilde{v}))_x = 0, \end{cases}$$

with the same initial data
 $(\tilde{u}, \tilde{v})(0, x)$

yield solutions are sufficiently close in the following sense:

- L^1 stability of $L^1 \cap BV$ solution with respect to flux functions, and
- Structural Stability of Riemann Problems; ie solutions with initial data of the form

$$(u, v)(0, x) = \begin{cases} (u_l, v_l) & \text{for } x < 0, \\ (u_r, v_r) & \text{for } x > 0. \end{cases}$$

Main Result

Consider:

$$\begin{cases} u_t + (F_0(u, v))_x = 0, \\ v_t + (G_0(u, v))_x = 0, \end{cases}$$

with Riemann initial data

$$(u, v)(0, x) = \begin{cases} (u_l, v_l) & \text{for } x < 0, \\ (u_r, v_r) & \text{for } x > 0, \end{cases}$$

for $(t, x) \in [0, \infty) \times \mathbb{R}$, $U \subset \mathbb{R}^2$ open, and $F_0, G_0 \in C^2(U)$.

Main Result

Assumptions: (F_0, G_0) forms a

(i) **Strictly hyperbolic** system in U :

Jacobian matrix $J(u, v; F_0, G_0) = \begin{pmatrix} (F_0)_u & (F_0)_v \\ (G_0)_u & (G_0)_v \end{pmatrix}$ possess two distinct real eigenvalues for each $(u, v) \in U$.

(ii) **Genuinely non-linear** system in U : For $k \in \{1, 2\}$

$\nabla \underbrace{\lambda_k}_{\text{k-Eigenvalue}}(u, v; F_0, G_0) \cdot \underbrace{\mathbf{r}_k}_{\text{k-Right Eigenvector}}(u, v; F_0, G_0) \neq 0$.

Convention: $\lambda_1 < \lambda_2$.

(iii) **Uni-directional system** in U : Either

- $(F_0)_v(u, v) \neq 0$ for all $(u, v) \in U$ or
- $(G_0)_u(u, v) \neq 0$ for all $(u, v) \in U$.

Main Result

Assumptions: (F_0, G_0) forms a

(i) **Strictly hyperbolic** system in U :

Jacobian matrix $J(u, v; F_0, G_0) = \begin{pmatrix} (F_0)_u & (F_0)_v \\ (G_0)_u & (G_0)_v \end{pmatrix}$ possess two distinct real eigenvalues for each $(u, v) \in U$.

(ii) **Genuinely non-linear** system in U : For $k \in \{1, 2\}$

$$\nabla \underbrace{\lambda_k}_{k\text{-Eigenvalue}}(u, v; F_0, G_0) \cdot \underbrace{\mathbf{r}_k}_{k\text{-Right Eigenvector}}(u, v; F_0, G_0) \neq 0.$$

Convention: $\lambda_1 < \lambda_2$.

(iii) **Uni-directional system** in U : Either

- $(F_0)_v(u, v) \neq 0$ for all $(u, v) \in U$ or
- $(G_0)_u(u, v) \neq 0$ for all $(u, v) \in U$.

Perturbations:

$$\begin{cases} \tilde{F} = F_0 + F_\delta, \\ \tilde{G} = G_0 + G_\delta. \end{cases}$$

Theorem (Generic Approximation Theorem & Structural Stability.)

Consider the system satisfying assumptions (i), (ii), (iii), and the regular manifold assumption. For almost every $(u_l, v_l) \neq (u_r, v_r) \in U$, consider the system with Riemann initial data such that there is a unique double-wave entropy solution. Then, for any compact subset $K \subset U$ containing (u_l, v_l) and (u_r, v_r) in its interior,

- 1 The unperturbed system satisfies the transversality property on K ,
- 2 There exists $\varepsilon_1, \varepsilon_2 > 0$ such that for any perturbations $(F_\delta, G_\delta) \in C^2(K)^2$ with $\|(F_\delta, G_\delta)\|_{C^2(K)^2} < \varepsilon_1$, the corresponding perturbed 2×2 system admits a unique double-wave entropy solution with an intermediate state $(\tilde{u}^*, \tilde{v}^*) \in \text{int}(K)$ satisfying $\|(\tilde{u}^*, \tilde{v}^*) - (u^*, v^*)\|_2 < \varepsilon_2$.
- 3 The perturbed system satisfies the transversality property on the same compact set K .

Main Result

Roughly speaking, this translates to:

For a system satisfying assumptions (i), (ii), (iii), and the regular manifold assumption, for almost every (generic) Riemann initial data, unique solutions and their structures (shock/rarefactions) are preserved upon a sufficiently small C^2 perturbation to the flux functions.

Furthermore, the “amplitudes” of shock and rarefaction upon perturbation are only perturbed by a small amount.


Crash Course: 2×2 System

$$\text{General System: } \begin{cases} u_t + (F(u, v))_x = 0, \\ v_t + (G(u, v))_x = 0. \end{cases}$$


Riemann initial data:

$$(u, v)(0, x) = \begin{cases} (u_l, v_l) & \text{for } x < 0, \\ (u_r, v_r) & \text{for } x > 0, \end{cases}$$

Shock:



Rarefaction:



Crash Course: 2×2 System

State Space (u, v) Analysis :

Given a state (u_l, v_l) ,

- (Shocks) Hugoniot loci: (Rankine-Hugoniot) All (u, v) satisfying $\begin{pmatrix} F(u, v) - F(u_l, v_l) \\ G(u, v) - G(u_l, v_l) \end{pmatrix} = s \begin{pmatrix} u - u_l \\ v - v_l \end{pmatrix}$ for some s .

Equivalently,

$$(F(u, v) - F(u_l, v_l))(v - v_l) - (G(u, v) - G(u_l, v_l))(u - u_l) = 0.$$

Required to satisfy 1-wave Lax Entropy condition.

- 1-Rarefaction Curves: All (u, v) solving

$$\begin{cases} \frac{d}{d\lambda}(u(\lambda), v(\lambda)) = \mathbf{r}_1(u(\lambda), v(\lambda)) \\ (u(\lambda(u_l, v_l)), v(\lambda(u_l, v_l))) = (u_l, v_l) \end{cases}$$

Integral curves of the right eigenvector \mathbf{r}_1 .

Solve for increasing λ .

Crash Course: 2×2 System

State Space (u, v) Analysis :

Given a state (u_r, v_r) ,

- (Shocks) Hugoniot loci: (Rankine-Hugoniot) All (u, v) satisfying $\begin{pmatrix} F(u, v) - F(u_r, v_r) \\ G(u, v) - G(u_r, v_r) \end{pmatrix} = s \begin{pmatrix} u - u_r \\ v - v_r \end{pmatrix}$ for some s .

Equivalently,

$$(F(u, v) - F(u_r, v_r))(v - v_r) - (G(u, v) - G(u_r, v_r))(u - u_r) = 0.$$

Required to satisfy 2-wave Lax Entropy condition.

- 2-Rarefaction Curves: All (u, v) solving

$$\begin{cases} \frac{d}{d\lambda}(u(\lambda), v(\lambda)) = \mathbf{r}_2(u(\lambda), v(\lambda)) \\ (u(\lambda(u_l, v_l)), v(\lambda(u_l, v_l))) = (u_l, v_l) \end{cases}$$


Integral curves of the right eigenvector \mathbf{r}_2 .

Solve for decreasing λ .

Crash Course: 2×2 System

Constructing Composite Solutions:

$$\underbrace{(u_l, v_l) \xrightarrow[\text{1-shock}]{\text{Hugoniot Locus}} (u^*, v^*) \xrightarrow[\text{2-rarefaction}]{\text{Rarefaction Curve}} (u_r, v_r)}_{\text{Shock-Rarefaction Solution}}$$




Hugoniot Loci /
Rarefaction Curve \

Crash Course: 2×2 System

Constructing Composite Solutions:

$$(u_l, v_l) \xrightarrow[\text{1-shock}]{\text{Hugoniot Locus}} (u^*, v^*) \xrightarrow[\text{2-shock}]{\text{Hugoniot Locus}} (u_r, v_r)$$


Double Shock



Hugoniot Loci /
Rarefaction Curve \

Unstable Case I: Single Wave Solution.



$$\underbrace{(u_l, v_l) \xrightarrow[\text{1-shock}]{\text{Hugoniot Locus}} (u^*, v^*) = (u_r, v_r)}_{\text{Single Shock}}$$



Hugoniot Loci /
 Rarefaction Curve ···

2×2 System

Unstable Case II: Self-Intersecting Hugoniot Loci.




Hugoniot Loci /
Rarefaction Curve ···


2×2 System

Unstable Case III: Singular (δ) Shock - Intersection at ∞ .
To be interpreted in the sense of distributions.
(Wang and Bertozzi, 2014.)

$$\underbrace{(u_l, v_l) \xrightarrow[\text{1-shock}]{\text{Hugoniot Locus}} \infty \xleftarrow[\text{2-shock}]{\text{Hugoniot Locus}} (u_r, v_r)}_{\text{Singular Shock}}$$




Hugoniot Loci /
Rarefaction Curve ···



2×2 System

Case IV: Singular Shock - Self-intersecting at given states.
(Keyfitz and Kranzer, 1990.)



Hugoniot Loci is **not** a manifold (locally ' \times ', not Euclidean).

Regular Manifold Assumption

Recall: Hugoniot loci connects all (u, v) from a given state (u_g, v_g)

$$(F(u, v) - F(u_g, v_g))(v - v_g) - (G(u, v) - G(u_g, v_g))(u - u_g) = 0.$$

Define the **Hugoniot Objective Function**:

$$H_{(u_g, v_g)} = (F(u, v) - F(u_g, v_g))(v - v_g) - (G(u, v) - G(u_g, v_g))(u - u_g).$$

Hugoniot locus is the zero level set of $H_{(u_g, v_g)}$.

Regular Manifold Assumption

Recall: Hugoniot loci connects all (u, v) from a given state (u_g, v_g)

$$(F(u, v) - F(u_g, v_g))(v - v_g) - (G(u, v) - G(u_g, v_g))(u - u_g) = 0.$$

Define the **Hugoniot Objective Function**:

$$H_{(u_g, v_g)} = (F(u, v) - F(u_g, v_g))(v - v_g) - (G(u, v) - G(u_g, v_g))(u - u_g).$$

Hugoniot locus is the zero level set of $H_{(u_g, v_g)}$.

Regular Manifold Assumption

The Jacobian map $(dH_{(u_g, v_g)})_{(u, v)} : \mathbb{R}^2 \rightarrow \mathbb{R}$ given by $(D_u H_{(u_g, v_g)}(u, v) \quad D_v H_{(u_g, v_g)}(u, v))$ is surjective for each $(u, v) \neq (u_g, v_g)$ on the Hugoniot locus.

- Always not satisfied at $(u, v) = (u_g, v_g)$.
- By the **Regular Value Theorem**, the Hugoniot locus restricted on $U \setminus \{(u_g, v_g)\}$ is a C^1 manifold.

Transversality


Let \mathcal{M} and \mathcal{N} be submanifolds of \mathbb{R}^n .

Definition: Transverse Intersection

We say that \mathcal{M} and \mathcal{N} **intersects transversely** if for every $x \in \mathcal{M} \cap \mathcal{N}$,

$$T_x\mathcal{M} + T_x\mathcal{N} = \mathbb{R}^n.$$

Notation: $\mathcal{M} \pitchfork \mathcal{N}$.




Transversality Property

Let K be a compact subset of U containing the given left and right states $(u_l, v_l) \neq (u_r, v_r)$.

Definition: Transversality Property

We say that the 2×2 system with Riemann initial data given by (u_l, v_l) and (u_r, v_r) as left and right states satisfies the **transversality property on K** if for the “correct” curves \mathcal{W}_l (from (u_l, v_l)) and \mathcal{W}_r (from (u_r, v_r)) intersecting at $(u^*, v^*) \neq (u_l, v_l)$ or (u_r, v_r) , we have

$$\mathcal{W}_l \pitchfork \mathcal{W}_r.$$



Theorem (Generic Approximation Theorem & Structural Stability.)

Consider the system satisfying assumptions (i), (ii), (iii), and the regular manifold assumption. For almost every $(u_l, v_l) \neq (u_r, v_r) \in U$, consider the system with Riemann initial data such that there is a unique double-wave entropy solution. Then, for any compact subset $K \subset U$ containing (u_l, v_l) and (u_r, v_r) in its interior,

- 1 The unperturbed system satisfies the transversality property on K ,
- 2 There exists $\varepsilon_1, \varepsilon_2 > 0$ such that for any perturbations $(F_\delta, G_\delta) \in C^2(K)^2$ with $\|(F_\delta, G_\delta)\|_{C^2(K)^2} < \varepsilon_1$, the corresponding perturbed 2×2 system admits a unique double-wave entropy solution with an intermediate state $(\tilde{u}^*, \tilde{v}^*) \in \text{int}(K)$ satisfying $\|(\tilde{u}^*, \tilde{v}^*) - (u^*, v^*)\|_2 < \varepsilon_2$.
- 3 The perturbed system satisfies the transversality property on the same compact set K .

Step I: Implicit Function Theorem on Banach Spaces

Theorem A. (Structural Stability)

Consider the system satisfying assumptions (i), (ii), (iii), and the regular manifold assumption. For **given states** $(u_l, v_l) \neq (u_r, v_r) \in U$, consider the system with Riemann initial data such that there is a unique double-wave entropy solution. Then, for any compact subset $K \subset U$ containing (u_l, v_l) and (u_r, v_r) in its interior,

If The unperturbed system satisfies the transversality property on K ,

Then There exists $\varepsilon_1, \varepsilon_2 > 0$ such that for any perturbations $(F_\delta, G_\delta) \in C^2(K)^2$ with $\|(F_\delta, G_\delta)\|_{C^2(K)^2} < \varepsilon_1$, the corresponding perturbed 2×2 system admits a unique double-wave entropy solution with an intermediate state $(\tilde{u}^*, \tilde{v}^*) \in \text{int}(K)$ satisfying $\|(\tilde{u}^*, \tilde{v}^*) - (u^*, v^*)\|_2 < \varepsilon_2$.

Moreover, The perturbed system satisfies the transversality property on the same compact set K .

Step I: Implicit Function Theorem on Banach Spaces

Proof Sketch (Persistence of Existence):

- **Hugoniot Objective Function** $H(u, v; u_g, v_g, F, G)$ given by

$$\begin{aligned} & H(u, v; u_g, v_g, F, G) \\ &= (F(u, v) - F(u_g, v_g))(v - v_g) - (G(u, v) - G(u_g, v_g))(u - u_g). \end{aligned}$$

- Hugoniot locus: All (u, v) such that $H(u, v; u_g, v_g, F, G) = 0$.

Step I: Implicit Function Theorem on Banach Spaces

Rarefaction Curves:

- Rarefaction ODEs:

$$\begin{cases} \frac{d}{d\lambda}(u(\lambda), v(\lambda)) = \mathbf{r}_k(u(\lambda), v(\lambda)) \\ (u(\lambda(u_g, v_g)), v(\lambda(u_g, v_g))) = (u_g, v_g) \end{cases}$$

- Use uni-direction assumption (iii) to normalize the 2nd component of the right eigenvector to be 1.
- Obtain a single ODE “ $\frac{du}{dv} = \frac{du/d\lambda}{dv/d\lambda}$ ”:

$$\begin{cases} \frac{d}{dv}u_k(v; F, G) = \Xi(u_k(v; F, G), F, G) \\ u_k(v_g) = u_g. \end{cases}$$

Step I: Implicit Function Theorem on Banach Spaces

Rarefaction Curves:

- Rarefaction ODEs:

$$\begin{cases} \frac{d}{d\lambda}(u(\lambda), v(\lambda)) = \mathbf{r}_k(u(\lambda), v(\lambda)) \\ (u(\lambda(u_g, v_g)), v(\lambda(u_g, v_g))) = (u_g, v_g) \end{cases}$$

- Use uni-direction assumption (iii) to normalize the 2nd component of the right eigenvector to be 1.
- Obtain a single ODE “ $\frac{du}{dv} = \frac{du/d\lambda}{dv/d\lambda}$ ”:

$$\begin{cases} \frac{d}{dv}u_k(v; F, G) = \Xi(u_k(v; F, G), F, G) \\ u_k(v_g) = u_g. \end{cases}$$

- **Rarefaction Objective Function:**


$$R_k(u, v; u_g, v_g, F, G) = u - u_k(v; u_g, v_g, F, G).$$

Step I: Implicit Function Theorem on Banach Spaces

Rarefaction Objective Function:

$$R_k(u, v; u_g, v_g, F, G) = u - u_k(v; u_g, v_g, F, G)$$

Interpretation: Signed Distance of u -coordinate to rarefaction curve integrated up to v .




Rarefaction Curve 

Step I: Implicit Function Theorem on Banach Spaces

Rarefaction Objective Function:

$$R_k(u, v; u_g, v_g, F, G) = u - u_k(v; u_g, v_g, F, G)$$


Interpretation: Signed Distance of u -coordinate to rarefaction curve integrated up to v .



Rarefaction Curve

k -Rarefaction curve = Zero-level set of R_k .

Step I: Implicit Function Theorem on Banach Spaces



Example: Unique intermediate state (u^*, v^*) and unperturbed fluxes (F_0, G_0) satisfy

$$\begin{cases} H(u^*, v^*, F_0, G_0; u_l, v_l) = 0, \\ R_2(u^*, v^*, F_0, G_0; u_r, v_r) = 0. \end{cases}$$

Step I: Implicit Function Theorem on Banach Spaces

Example: Unique intermediate state (u^*, v^*) and unperturbed fluxes (F_0, G_0) satisfy

$$\begin{cases} H(u^*, v^*, F_0, G_0; u_l, v_l) = 0, \\ R_2(u^*, v^*, F_0, G_0; u_r, v_r) = 0. \end{cases}$$

Step I: Implicit Function Theorem on Banach Spaces

Example: Unique intermediate state (u^*, v^*) and unperturbed fluxes (F_0, G_0) satisfy

$$\begin{cases} H(u^*, v^*, F_0, G_0; u_l, v_l) = 0, \\ R_2(u^*, v^*, F_0, G_0; u_r, v_r) = 0. \end{cases}$$

Apply **Implicit Function Theorem on Banach Spaces** to

$$\mathbf{hr}(u, v, F, G) := \begin{cases} H(\mathbf{u}, \mathbf{v}, \mathbf{F}, \mathbf{G}; u_l, v_l) = 0, \\ R_2(\mathbf{u}, \mathbf{v}, \mathbf{F}, \mathbf{G}; u_r, v_r) = 0. \end{cases}$$

with $(u, v) \in K$ and $(F, G) \in C^2(K)^2$ to obtain a map $M : C^2(K)^2 \rightarrow K$ such that

Step I: Implicit Function Theorem on Banach Spaces

Example: Unique intermediate state (u^*, v^*) and unperturbed fluxes (F_0, G_0) satisfy

$$\begin{cases} H(u^*, v^*, F_0, G_0; u_l, v_l) = 0, \\ R_2(u^*, v^*, F_0, G_0; u_r, v_r) = 0. \end{cases}$$

Apply **Implicit Function Theorem on Banach Spaces** to


$$\mathbf{hr}(u, v, F, G) := \begin{cases} H(\mathbf{u}, \mathbf{v}, \mathbf{F}, \mathbf{G}; u_l, v_l) = 0, \\ R_2(\mathbf{u}, \mathbf{v}, \mathbf{F}, \mathbf{G}; u_r, v_r) = 0. \end{cases}$$

with $(u, v) \in K$ and $(F, G) \in C^2(K)^2$ to obtain a map $M : C^2(K)^2 \rightarrow K$ such that

$$\begin{cases} H(M(F, G), F, G; u_l, v_l) = 0, \\ R_2(M(F, G), F, G; u_r, v_r) = 0. \end{cases}$$

with $M(F_0, G_0) = (u^*, v^*)$ in a $C^2(K)^2$ neighborhood of (F_0, G_0) .

Step I: Implicit Function Theorem on Banach Spaces



Transition to Step II

$$\begin{cases} H(M(F, G), F, G; u_l, v_l) = 0, \\ R_2(M(F, G), F, G; u_r, v_r) = 0. \end{cases}$$

Transition to Step II

$$\begin{cases} H(M(F, G), F, G; u_l, v_l) = 0, \\ R_2(M(F, G), F, G; u_r, v_r) = 0. \end{cases}$$

To apply **Implicit Function Theorem on Banach Spaces**, we need to check the invertibility of a Jacobian map (matrix)

$D_{(u,v)} \mathbf{hr}(u^*, v^*, F_0, G_0) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, given by

$$\begin{pmatrix} D_u H(u^*, v^*, F_0, G_0; u_l, v_l) & D_v H(u^*, v^*, F_0, G_0; u_l, v_l) \\ D_u R_2(u^*, v^*, F_0, G_0; u_r, v_r) & D_v R_2(u^*, v^*, F_0, G_0; u_r, v_r) \end{pmatrix}.$$

Transition to Step II

$$\begin{cases} H(M(F, G), F, G; u_l, v_l) = 0, \\ R_2(M(F, G), F, G; u_r, v_r) = 0. \end{cases}$$

To apply **Implicit Function Theorem on Banach Spaces**, we need to check the invertibility of a Jacobian map (matrix)

$D_{(u,v)} \mathbf{hr}(u^*, v^*, F_0, G_0) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, given by

$$\begin{pmatrix} D_u H(u^*, v^*, F_0, G_0; u_l, v_l) & D_v H(u^*, v^*, F_0, G_0; u_l, v_l) \\ D_u R_2(u^*, v^*, F_0, G_0; u_r, v_r) & D_v R_2(u^*, v^*, F_0, G_0; u_r, v_r) \end{pmatrix}.$$

If $\mathcal{W}_l =$ Hugoniot locus from (u_l, v_l) and $\mathcal{W}_r =$ Rarefaction curve from (u_r, v_r) , this is equivalent to

$$\mathcal{W}_l \pitchfork \mathcal{W}_r = \mathbb{R}^2.$$

Transition to Step II

Recall:

Theorem A. (Structural Stability)

Consider the system satisfying assumptions (i), (ii), (iii), and the regular manifold assumption. For **given states** $(u_l, v_l) \neq (u_r, v_r) \in U$, consider the system with Riemann initial data such that there is a unique double-wave entropy solution. Then, for any compact subset $K \subset U$ containing (u_l, v_l) and (u_r, v_r) in its interior,

If The unperturbed system satisfies the transversality property on K ,

Then There exists $\varepsilon_1, \varepsilon_2 > 0$ such that for any perturbations $(F_\delta, G_\delta) \in C^2(K)^2$ with $\|(F_\delta, G_\delta)\|_{C^2(K)^2} < \varepsilon_1$, the corresponding perturbed 2×2 system admits a unique double-wave entropy solution with an intermediate state $(\tilde{u}^*, \tilde{v}^*) \in \text{int}(K)$ satisfying $\|(\tilde{u}^*, \tilde{v}^*) - (u^*, v^*)\|_2 < \varepsilon_2$.

Moreover, The perturbed system satisfies the transversality property on the same compact set K .

Transition to Step II

Theorem B. (Transversality is Generic.)

Consider the system satisfying assumptions (i), (ii), (iii), and the regular manifold assumption. For **almost every** $(u_l, v_l) \neq (u_r, v_r) \in U$, consider the system with Riemann initial data such that there is a unique double-wave entropy solution. Then, for any compact subset $K \subset U$ containing (u_l, v_l) and (u_r, v_r) in its interior, we

Get: The unperturbed system satisfies the transversality property on K .

Transition to Step II

Theorem A + Theorem B = Main Theorem.

Theorem. (Generic Approximation Theorem & Structural Stability.)

Consider the system satisfying assumptions (i), (ii), (iii), and the regular manifold assumption. For **almost every** $(u_l, v_l) \neq (u_r, v_r) \in U$, consider the system with Riemann initial data such that there is a unique double-wave entropy solution. Then, for any compact subset $K \subset U$ containing (u_l, v_l) and (u_r, v_r) in its interior,

- 1 The unperturbed system satisfies the transversality property on K ,
- 2 There exists $\varepsilon_1, \varepsilon_2 > 0$ such that for any perturbations $(F_\delta, G_\delta) \in C^2(K)^2$ with $\|(F_\delta, G_\delta)\|_{C^2(K)^2} < \varepsilon_1$, the corresponding perturbed 2×2 system admits a unique double-wave entropy solution with an intermediate state $(\tilde{u}^*, \tilde{v}^*) \in \text{int}(K)$ satisfying $\|(\tilde{u}^*, \tilde{v}^*) - (u^*, v^*)\|_2 < \varepsilon_2$.
- 3 The perturbed system satisfies the transversality property on the same compact set K .

Step II: Parametric Transversality Theorems

Theorem B. (Transversality is Generic.)

Consider the system satisfying assumptions (i), (ii), (iii), and the regular manifold assumption. For **almost every** $(u_l, v_l) \neq (u_r, v_r) \in U$, consider the system with Riemann initial data such that there is a unique double-wave entropy solution. Then, for any compact subset $K \subset U$ containing (u_l, v_l) and (u_r, v_r) in its interior, we

Get: The unperturbed system satisfies the transversality property on K .

Step II: Parametric Transversality Theorems

Let \mathcal{X} and \mathcal{Y} be C^r manifolds, and \mathcal{Z} be a C^r submanifold of \mathcal{Y} for $r \geq 1$.


Definition: Transversality of a Map

Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a C^r map. We say that f is **transverse** to \mathcal{Z} if for every $a \in f^{-1}(\mathcal{Z})$, we have

$$df(T_a\mathcal{X}) + T_{f(a)}\mathcal{Z} = T_{f(a)}\mathcal{Y}.$$

Notation: $f \pitchfork \mathcal{Z}$.

Intuition: " $f(\mathcal{X}) \pitchfork \mathcal{Z}$ ".



Step II: Parametric Transversality Theorems

Typical genericity arguments utilize:

Thom's Parametric Transversality Theorem

Let \mathcal{X} , \mathcal{P} , and \mathcal{Y} be C^r manifolds and \mathcal{Z} be a C^r submanifolds of \mathcal{N} . Consider


- The map $F : \mathcal{X} \times \mathcal{P} \rightarrow \mathcal{Y}$, and
- The associated parametric maps $F_p : \mathcal{X} \rightarrow \mathcal{Y}$ for each $p \in \mathcal{P}$.

Suppose that

- 1 $r > \max\{0, \dim \mathcal{Y} + \dim \mathcal{Z} - \dim \mathcal{X}\}$,
- 2 The map $(x, p) \mapsto F_p(x)$ is C^r , and
- 3 $F \pitchfork \mathcal{Z}$.

Then, for almost every $p \in \mathcal{P}$, $F_p \pitchfork \mathcal{Z}$.


Step II: Parametric Transversality Theorems



Strategy 1: $\mathbf{hr}(\cdot; u_l, v_l, u_r, v_r) : U \rightarrow \mathbb{R}^2$ with

$$\mathbf{hr}(u, v; u_l, v_l, u_r, v_r) := \begin{cases} H(u, v; u_l, v_l), \\ R_2(u, v; u_r, v_r). \end{cases}$$

Step II: Parametric Transversality Theorems




Strategy 1: $\mathbf{hr}(\cdot; u_l, v_l, u_r, v_r) : U \rightarrow \mathbb{R}^2$ with

$$\mathbf{hr}(u, v; u_l, v_l, u_r, v_r) := \begin{cases} H(u, v; u_l, v_l), \\ R_2(u, v; u_r, v_r). \end{cases}$$

Recall: $\mathbf{hr}(u^*, v^*; u_l, v_l, u_r, v_r) = \mathbf{0}$.

Step II: Parametric Transversality Theorems




Strategy 1: $\mathbf{hr}(\cdot; u_l, v_l, u_r, v_r) : U \rightarrow \mathbb{R}^2$ with

$$\mathbf{hr}(u, v; u_l, v_l, u_r, v_r) := \begin{cases} H(u, v; u_l, v_l), \\ R_2(u, v; u_r, v_r). \end{cases}$$

Recall: $\mathbf{hr}(u^*, v^*; u_l, v_l, u_r, v_r) = \mathbf{0}$.

Hope: $\mathbf{hr}(\cdot; u_l, v_l, u_r, v_r) \pitchfork \{(0, 0)\}$ for almost every $(u_l, v_l) \neq (u_r, v_r)$.

Step II: Parametric Transversality Theorems



Strategy 1: $\mathbf{hr}(\cdot; u_l, v_l, u_r, v_r) : U \rightarrow \mathbb{R}^2$ with

$$\mathbf{hr}(u, v; u_l, v_l, u_r, v_r) := \begin{cases} H(u, v; u_l, v_l), \\ R_2(u, v; u_r, v_r). \end{cases}$$

Recall: $\mathbf{hr}(u^*, v^*; u_l, v_l, u_r, v_r) = \mathbf{0}$.

Hope: $\mathbf{hr}(\cdot; u_l, v_l, u_r, v_r) \pitchfork \{(0, 0)\}$ for almost every $(u_l, v_l) \neq (u_r, v_r)$.

Conclude: At each intersection point \implies transverse intersection.

Step II: Parametric Transversality Theorems

Define $\Delta_{U^2} = \{(u_l, v_l, u_r, v_r) : (u_l, v_l) \neq (u_r, v_r)\}$.

To apply **Thom's Parametric Transversality Theorem**, we need to check that

$$\mathbf{hr} : \underbrace{U}_{\text{Allowed Intersection Points}} \times \underbrace{(U^2 \setminus \Delta_{U^2})}_{\text{Parameters: Left and Right States}} \rightarrow \mathbb{R}^2$$

satisfies $\mathbf{hr} \pitchfork \{(0, 0)\}$.

Step II: Parametric Transversality Theorems

Define $\Delta_{U^2} = \{(u_l, v_l, u_r, v_r) : (u_l, v_l) \neq (u_r, v_r)\}$.

To apply **Thom's Parametric Transversality Theorem**, we need to check that

$$\mathbf{hr} : \underbrace{U}_{\text{Allowed Intersection Points}} \times \underbrace{(U^2 \setminus \Delta_{U^2})}_{\text{Parameters: Left and Right States}} \rightarrow \mathbb{R}^2$$

satisfies $\mathbf{hr} \pitchfork \{(0, 0)\}$.

Then, $\mathbf{hr}(\cdot; u_l, v_l, u_r, v_r) \pitchfork \{(0, 0)\}$ for almost every $(u_l, v_l, u_r, v_r) \in U^2 \setminus \Delta_{U^2}$.

Step II: Parametric Transversality Theorems

Define $\Delta_{U^2} = \{(u_l, v_l, u_r, v_r) : (u_l, v_l) \neq (u_r, v_r)\}$.

To apply **Thom's Parametric Transversality Theorem**, we need to check that

$$\mathbf{hr} : \underbrace{U}_{\text{Allowed Intersection Points}} \times \underbrace{(U^2 \setminus \Delta_{U^2})}_{\text{Parameters: Left and Right States}} \rightarrow \mathbb{R}^2$$

satisfies $\mathbf{hr} \pitchfork \{(0, 0)\}$.

Then, $\mathbf{hr}(\cdot; u_l, v_l, u_r, v_r) \pitchfork \{(0, 0)\}$ for almost every $(u_l, v_l, u_r, v_r) \in U^2 \setminus \Delta_{U^2}$.



Hugoniot Loci from (u_g, v_g) are manifolds on $U \setminus (u_g, v_g)$.
(i.e Keyfitz-Kranzer system.)

Step II: Parametric Transversality Theorems

Strategy 2: Puncture the domain U at (u_l, v_l) and (u_r, v_r) for **each** given left and right states.

Step II: Parametric Transversality Theorems

Strategy 2: Puncture the domain U at (u_l, v_l) and (u_r, v_r) for **each** given left and right states.

Set $U_{(u_l, v_l, u_r, v_r)} := U \setminus \{(u_l, v_l), (u_r, v_r)\}$ and define

$$ULR = \bigcup_{(u_l, v_l, u_r, v_r) \in U^2 \setminus \Delta_{U^2}} \underbrace{U_{(u_l, v_l, u_r, v_r)}}_{\text{Intersection Points}} \times \underbrace{\{(u_l, v_l, u_r, v_r)\}}_{\text{Parameters}}.$$

Step II: Parametric Transversality Theorems

Strategy 2: Puncture the domain U at (u_l, v_l) and (u_r, v_r) for **each** given left and right states.


Set $U_{(u_l, v_l, u_r, v_r)} := U \setminus \{(u_l, v_l), (u_r, v_r)\}$ and define

$$ULR = \bigcup_{(u_l, v_l, u_r, v_r) \in U^2 \setminus \Delta_{U^2}} \underbrace{U_{(u_l, v_l, u_r, v_r)}}_{\text{Intersection Points}} \times \underbrace{\{(u_l, v_l, u_r, v_r)\}}_{\text{Parameters}}.$$

Now, check $\mathbf{hr} : ULR \rightarrow \mathbb{R}^2$ satisfies

$$\mathbf{hr} \pitchfork \{(0, 0)\}.$$

Step II: Parametric Transversality Theorems



$$ULR = \bigcup_{(u_l, v_l, u_r, v_r) \in U^2 \setminus \Delta_{U^2}} U(u_l, v_l, u_r, v_r) \times \{(u_l, v_l, u_r, v_r)\}.$$

ULR is a 6-dimensional submanifold of \mathbb{R}^6 .

Step II: Parametric Transversality Theorems

Foliated Parametric Transversality Theorem

Let \mathcal{P} and \mathcal{Y} be C^r manifolds, and \mathcal{Z} be a C^r submanifold of \mathcal{Y} . Suppose that for each $p \in \mathcal{P}$, we consider a collection of C^r manifolds given by $\{\mathcal{X}_p\}_{p \in \mathcal{P}}$ each with the same dimension $\dim \mathcal{X}$, and the following foliated set:



$$\mathcal{X}\mathcal{P} := \bigcup_{p \in \mathcal{P}} \mathcal{X}_p \times \{p\}. \quad (1)$$

Consider the maps $F : \mathcal{X}\mathcal{P} \rightarrow \mathcal{Y}$ and the associated map $F_p : \mathcal{X}_p \rightarrow \mathcal{Y}$ for each parameter $p \in \mathcal{P}$. Suppose that

1. $r > \max\{0, \dim \mathcal{Y} + \dim \mathcal{Z} - \dim \mathcal{X}\}$,
2. $\mathcal{X}\mathcal{P}$ is a C^r manifold with dimension $\dim \mathcal{X}\mathcal{P} = \dim \mathcal{X} + \dim \mathcal{P}$,
3. $T_{(x,p)}\mathcal{X}\mathcal{P} = T_x\mathcal{X}_p \times T_p\mathcal{P}$ for each $(x,p) \in \mathcal{X}\mathcal{P}$,
4. The map $(x,p) \mapsto F_p(x)$ is C^r , and
5. $F \pitchfork \mathcal{Z}$.

Then, for almost every $p \in \mathcal{P}$, $F_p \pitchfork \mathcal{Z}$.

Step II: Parametric Transversality Theorems



Theorem (Generic Approximation Theorem & Structural Stability.)

Consider the system satisfying assumptions (i), (ii), (iii), and the regular manifold assumption. For almost every $(u_l, v_l) \neq (u_r, v_r) \in U$, consider the system with Riemann initial data such that there is a unique double-wave entropy solution. Then, for any compact subset $K \subset U$ containing (u_l, v_l) and (u_r, v_r) in its interior,

- 1 The unperturbed system satisfies the transversality property on K ,
- 2 There exists $\varepsilon_1, \varepsilon_2 > 0$ such that for any perturbations $(F_\delta, G_\delta) \in C^2(K)^2$ with $\|(F_\delta, G_\delta)\|_{C^2(K)^2} < \varepsilon_1$, the corresponding perturbed 2×2 system admits a unique double-wave entropy solution with an intermediate state $(\tilde{u}^*, \tilde{v}^*) \in \text{int}(K)$ satisfying $\|(\tilde{u}^*, \tilde{v}^*) - (u^*, v^*)\|_2 < \varepsilon_2$.
- 3 The perturbed system satisfies the transversality property on the same compact set K .

$\mathbf{hr}(\underbrace{u, v}_{\text{Double-Wave Solutions}}; \underbrace{u_l, v_l, u_r, v_r}_{\text{Generically}}, \underbrace{F, G}_{\text{Perturbation}} \rightarrow \text{Stable})$

L^1 Stability:

L^1 Stability:

- (Holden and Holden, 1992.) L^1 stability for scalar conservation laws:

$$\|u_f(t, \cdot) - u_g(t, \cdot)\|_{L^1} \lesssim t \text{Lip}(f - g).$$

Done using the front-tracking algorithm.

L^1 Stability:

- (Holden and Holden, 1992.) L^1 stability for scalar conservation laws:

$$\|u_f(t, \cdot) - u_g(t, \cdot)\|_{L^1} \lesssim t \text{Lip}(f - g).$$

Done using the front-tracking algorithm.

- (Bianchini and Colombo, 2002.) L^1 stability for systems:

$$\left\| \begin{pmatrix} u \\ v \end{pmatrix} (t, \cdot) - \begin{pmatrix} \tilde{u} \\ \tilde{v} \end{pmatrix} (t, \cdot) \right\|_{L^1} \lesssim C_{(F_0, G_0)} C_{(\tilde{F}, \tilde{G})} \hat{d}((F_0, G_0), (\tilde{F}, \tilde{G})).$$

Done using (semi-)standard PDE techniques on Riemann semigroup.

Structural Stability of Riemann Problem:

- (Schechter, Marchesin, and Plohr, 1994.)
Structurally Stable Riemann Solutions.
 - Conclusion depends on the given left and right states and transversality condition of intersecting curves that could not be checked *a priori*.
 - Done using viscous regularization, traveling waves, and phase portrait analysis.

Structural Stability of Riemann Problem:

- (Schechter, Marchesin, and Plohr, 1994.)
Structurally Stable Riemann Solutions.
 - Conclusion depends on the given left and right states and transversality condition of intersecting curves that could not be checked *a priori*.
 - Done using viscous regularization, traveling waves, and phase portrait analysis.
- (Azevedo et. al., 2010 and Eschenazi et. al., 2025.)
Topological Approach for 2×2 systems.
 - Quadratic flux and perturbations; some work in progress.
 - Similar issue with transversality condition.
 - Done by employing desingularization methods (motivated by singularity theorem) specific to quadratic fluxes.

Genericity for Conservation Laws:

- (Schaeffer, 1973.)

Schaeffer Regularity Theorem (for scalar conservation laws): For almost any $u(0, x) \in \mathcal{S}(\mathbb{R})$, the solution is piecewise smooth with a finite number of shock curves.

- Only for scalar conservation laws; strong assumptions on initial data.
- Done using a transversality argument (differential topology).

Genericity for Conservation Laws:

- (Schaeffer, 1973.)
Schaeffer Regularity Theorem (for scalar conservation laws): For almost any $u(0, x) \in \mathcal{S}(\mathbb{R})$, the solution is piecewise smooth with a finite number of shock curves.
 - Only for scalar conservation laws; strong assumptions on initial data.
 - Done using a transversality argument (differential topology).
- (Caravenna and Spinolo, 2017.)
Schaeffer's Regularity Theorem Does Not Extend to Systems.

Genericity for Conservation Laws:

- (Schaeffer, 1973.)
Schaeffer Regularity Theorem (for scalar conservation laws): For almost any $u(0, x) \in \mathcal{S}(\mathbb{R})$, the solution is piecewise smooth with a finite number of shock curves.
 - Only for scalar conservation laws; strong assumptions on initial data.
 - Done using a transversality argument (differential topology).
- (Caravenna and Spinolo, 2017.)
Schaeffer's Regularity Theorem Does Not Extend to Systems.
- (Bressan, Chen, and Huang, 2024.)
Generic Singularities for 2D Pressureless Flow.
 - $x \in \mathbb{R}^2$, only for smooth initial data and a specific problem.

Application I: p-system

(Wendroff, 1972 and Wendroff, 1972.)

$$\begin{cases} u_t + (p(v))_x = 0, \\ v_t - u_x = 0. \end{cases}$$

Application I: p-system

(Wendroff, 1972 and Wendroff, 1972.)

$$\begin{cases} u_t + (p(v))_x = 0, \\ v_t - u_x = 0. \end{cases}$$

Compressible Isentropic Flow in Lagrangian Coordinates:

- Lagrangian Coordinates x
- Velocity in Lagrangian Coordinates $u \in \mathbb{R}$
- Specific Volume $v > 0$
- Pressure $p(v) \in C^2((0, \infty))$

Application I: p-system

(Wendroff, 1972 and Wendroff, 1972.)

$$\begin{cases} u_t + (p(v))_x = 0, \\ v_t - u_x = 0. \end{cases}$$

Compressible Isentropic Flow in Lagrangian Coordinates:

- Lagrangian Coordinates x
- Velocity in Lagrangian Coordinates $u \in \mathbb{R}$
- Specific Volume $v > 0$
- Pressure $p(v) \in C^2((0, \infty))$

Modelling Assumptions:

- Thermodynamics : $p'(v) < 0$ for $v > 0$.
- Experimental Evidence (Bethe, 1942): $p''(v) > 0$ for $v > 0$.

Application I: p-system

Jacobian Matrix:

$$J(u, v) = \begin{pmatrix} 0 & p'(v) \\ -1 & 0 \end{pmatrix}$$

Application I: p-system

Jacobian Matrix:

$$J(u, v) = \begin{pmatrix} 0 & p'(v) \\ -1 & 0 \end{pmatrix}$$

Assumptions:

- (i) $p'(v) < 0$ for $v > 0$ implies **strictly hyperbolic** system in $(0, \infty) \times \mathbb{R}$.
- (ii) $p''(v) > 0$ for $v > 0$ implies **genuinely non-linear** system in $(0, \infty) \times \mathbb{R}$.
- (iii) $-1 \neq 0$ implies **uni-directional system** in $(0, \infty) \times \mathbb{R}$.

Application I: p-system

Manifold Assumption:

Application I: p-system

Manifold Assumption:

- Hugoniot Objective Function:

$$H(u, v; u_g, v_g) = (u - u_g)^2 + (p(v) - p(v_g)) \cdot (v - v_g).$$

Application I: p-system

Manifold Assumption:

- Hugoniot Objective Function:

$$H(u, v; u_g, v_g) = (u - u_g)^2 + (p(v) - p(v_g)) \cdot (v - v_g).$$

- Jacobian:

$$(dH_{(u_g, v_g)})_{(u, v)} = (2(u - u_g) \quad p'(v)(v - v_g) + (p(v) - p(v_g))).$$

Application I: p-system

Manifold Assumption:

- Hugoniot Objective Function:

$$H(u, v; u_g, v_g) = (u - u_g)^2 + (p(v) - p(v_g)) \cdot (v - v_g).$$

- Jacobian:

$$(dH_{(u_g, v_g)})_{(u, v)} = (2(u - u_g) \quad p'(v)(v - v_g) + (p(v) - p(v_g))).$$

- Show that $(dH_{(u_g, v_g)})_{(u, v)} : \mathbb{R}^2 \rightarrow \mathbb{R}$ is surjective for any $(u, v) \neq (u_g, v_g)$ on the Hugoniot locus.

Application I: p-system

Manifold Assumption:

- Hugoniot Objective Function:

$$H(u, v; u_g, v_g) = (u - u_g)^2 + (p(v) - p(v_g)) \cdot (v - v_g).$$

- Jacobian:

$$(dH_{(u_g, v_g)})_{(u, v)} = (2(u - u_g) \quad p'(v)(v - v_g) + (p(v) - p(v_g))).$$

- Show that $(dH_{(u_g, v_g)})_{(u, v)} : \mathbb{R}^2 \rightarrow \mathbb{R}$ is surjective for any $(u, v) \neq (u_g, v_g)$ on the Hugoniot locus.

Physical Interpretation:

For a sufficiently good C^2 approximation of the pressure relation (on a compact subset), unique double-wave entropy solutions are preserved.

Application II: Gravity-Driven Particle-Laden Flow

$$\left\{ \begin{array}{l} h_t + \underbrace{\left(h^3 f \left(\frac{h\phi_0}{h} \right) \right)}_{F(h,h\phi_0)} \Big|_x = 0, \\ (h\phi_0)_t + \underbrace{\left(h^3 g \left(\frac{h\phi_0}{h} \right) \right)}_{G(h,h\phi_0)} \Big|_x = 0. \end{array} \right.$$

- $f(\phi_0), g(\phi_0), \phi_0 \in [0, \phi_m]$.

Physical Interpretation: $\phi_m =$ Maximum Packing Fraction.

Application II: Gravity-Driven Particle-Laden Flow

$$\left\{ \begin{array}{l} h_t + \underbrace{\left(h^3 f \left(\frac{h\phi_0}{h} \right) \right)}_{F(h, h\phi_0)} \Big|_x = 0, \\ (h\phi_0)_t + \underbrace{\left(h^3 g \left(\frac{h\phi_0}{h} \right) \right)}_{G(h, h\phi_0)} \Big|_x = 0. \end{array} \right.$$

- $f(\phi_0), g(\phi_0), \phi_0 \in [0, \phi_m]$.
Physical Interpretation: $\phi_m =$ Maximum Packing Fraction.
- If the assumptions of the theorem hold, by picking a sufficiently good approximation of f and g , hence F and G , unique double-wave entropy solutions are preserved.

Application II-1: Interpolating Flux Functions

Algorithm:

- 1 Place a grid on $\phi_0 = [0, \phi_m]$ with $\phi_m = 0.610$, say step size $\Delta\phi_0 = 0.001$.
- 2 Solve the nonlinear ODE for $\phi_0 = 0.001i$ for $i = 1, \dots, 610$ to obtain $f(\phi_0)$ and $g(\phi_0)$.
- 3 Obtain $f(\phi_0)$ and $g(\phi_0)$ by interpolation.
- 4 Obtain $f'(\phi_0)$ and $g'(\phi_0)$ by interpolation too (if needed).

Global Error for $f = \|f - f_{\text{int}}\|_{C^1(K)} \lesssim o(\Delta\phi_0)$

$$\xrightarrow{\Delta\phi_0 \rightarrow 0} 0$$

Application II-1: Interpolating Flux Functions

Algorithm:

- 1 Place a grid on $\phi_0 = [0, \phi_m]$ with $\phi_m = 0.610$, say step size $\Delta\phi_0 = 0.001$.
- 2 Solve the nonlinear ODE for $\phi_0 = 0.001i$ for $i = 1, \dots, 610$ to obtain $f(\phi_0)$ and $g(\phi_0)$.
- 3 Obtain $f(\phi_0)$ and $g(\phi_0)$ by interpolation.
- 4 Obtain $f'(\phi_0)$ and $g'(\phi_0)$ by interpolation too (if needed).

$$\text{Global Error for } f = \|f - f_{\text{int}}\|_{C^1(K)} \lesssim o(\Delta\phi_0)$$
$$\xrightarrow{\Delta\phi_0 \rightarrow 0} 0$$

Interpretation:

The solutions exhibit structural stability for a sufficiently small grid size, with solutions converging to the original system as grid size goes to 0.

Application II-2: Gravity-Driven Particle-Laden Flow


Fix $\alpha = 25^\circ$. $\phi_0 \in [0, \phi_m]$,

$\phi_m = 0.61$: Maximum packing fraction.

$\phi_c \approx 0.503$: Phase transition from settled to ridged.

Settled: $\phi_0 < \phi_c$.

Ridged: $\phi_0 > \phi_c$.



Application II-2: Gravity-Driven Particle-Laden Flow

Fix $\alpha = 25^\circ$. $\phi_0 \in [0, \phi_m]$,

$\phi_m = 0.61$: Maximum packing fraction.

$\phi_c \approx 0.503$: Phase transition from settled to ridged.

Settled: $\phi_0 < \phi_c$.

Ridged: $\phi_0 > \phi_c$.



Polynomial Approximations:

$$f(\phi_0) = \begin{cases} \sum_{j=1}^{10} \beta_{f,j}^S (\phi_c - \phi_0)^{j-1} & \text{for } \phi_0 < \phi_c, \\ \sum_{j=1}^{10} \beta_{f,j}^R (\phi_0 - \phi_c)^{j-1} & \text{for } \phi_0 > \phi_c, \end{cases}$$

and

$$g(\phi_0) = \begin{cases} \sum_{j=1}^{10} \beta_{g,j}^S (\phi_c - \phi_0)^{j-1} & \text{for } \phi_0 < \phi_c, \\ \sum_{j=1}^{10} \beta_{g,j}^R (\phi_0 - \phi_c)^{j-1} & \text{for } \phi_0 > \phi_c. \end{cases}$$

$$\beta_f = \operatorname{argmin}_{\beta_f \in \mathbb{R}^{20}} \|\mathbf{f} - \mathbf{X}\beta_f\|_2^2 + \lambda \|\mathbf{f}' - \mathbf{X}'\beta_f\|_2^2$$

subject to the assumptions above (similar for g).

Physical “Constraints”:

- (I) : $f, f', f'', g, g',$ and g'' are continuous at ϕ_c ,
- (II) : $f(0) = \frac{\mu l}{3}, g(0) = 0,$
- (III) : Values of $f(\phi_c)$ and $g(\phi_c),$
- (IV) : $f(\phi_m) = g(\phi_m) = f'(\phi_m) = g'(\phi_m) = 0.$

$$\beta_f = \operatorname{argmin}_{\beta_f \in \mathbb{R}^{20}} \|\mathbf{f} - \mathbf{X}\beta_f\|_2^2 + \lambda \|\mathbf{f}' - \mathbf{X}'\beta_f\|_2^2$$

subject to the assumptions above (similar for g).

Physical “Constraints”:

- (I) : f, f', f'', g, g' , and g'' are continuous at ϕ_c ,
- (II) : $f(0) = \frac{\mu l}{3}, g(0) = 0$,
- (III) : Values of $f(\phi_c)$ and $g(\phi_c)$,
- (IV) : $f(\phi_m) = g(\phi_m) = f'(\phi_m) = g'(\phi_m) = 0$.

Compare:

- $\lambda = 0$ (not fitting for derivatives) and
- $\lambda = 0.03$ (fitting for derivatives, obtained via leave-one-out cross validation).

$$\beta_f = \operatorname{argmin}_{\beta_f \in \mathbb{R}^{20}} \|\mathbf{f} - \mathbf{X}\beta_f\|_2^2 + \lambda \|\mathbf{f}' - \mathbf{X}'\beta_f\|_2^2$$

subject to the assumptions above (similar for g). Sampled Data

Points:

- A couple of points close to ϕ_m ,
- A couple of points close to ϕ_c ,
- A couple of sparse points,
- Points are in triplets to provide derivative information at the middle point.

Optimization Algorithm:

- Quadratic program with linear equality constraints.
- Determine λ by using a leave-one-out cross validation algorithm.

Order of Phase Transition

C^1 vs C^2 ?

Order of Phase Transition

C^1 vs C^2 ?

Asymptotically,

$$f(\phi_0), g(\phi_0) \sim |\phi_0 - \phi_c|^\beta$$

with

- $\beta < 1$ if $\alpha > 70.309^\circ$
- $\beta \in (1, 2)$ if $\alpha \in (27.895^\circ, 70.309^\circ)$
- $\beta > 2$ if $\alpha < 27.895^\circ$

Order of Phase Transition

C^1 vs C^2 ?

Asymptotically,

$$f(\phi_0), g(\phi_0) \sim |\phi_0 - \phi_c|^\beta$$

with

- $\beta < 1$ if $\alpha > 70.309^\circ$
- $\beta \in (1, 2)$ if $\alpha \in (27.895^\circ, 70.309^\circ)$
- $\beta > 2$ if $\alpha < 27.895^\circ$

It can be “numerically verified” that $f(\phi_0)$ and $g(\phi_0)$ are

- C^2 across $\phi_0 = \phi_c$ for $\alpha = 17^\circ$.
- C^1 only across $\phi_0 = \phi_c$ for $\alpha = 30^\circ, 60^\circ, 80^\circ$.

Order of Phase Transition

C^1 vs C^2 ?

Asymptotically,

$$f(\phi_0), g(\phi_0) \sim |\phi_0 - \phi_c|^\beta$$

with

- $\beta < 1$ if $\alpha > 70.309^\circ$
- $\beta \in (1, 2)$ if $\alpha \in (27.895^\circ, 70.309^\circ)$
- $\beta > 2$ if $\alpha < 27.895^\circ$


It can be “numerically verified” that $f(\phi_0)$ and $g(\phi_0)$ are

- C^2 across $\phi_0 = \phi_c$ for $\alpha = 17^\circ$.
- C^1 only across $\phi_0 = \phi_c$ for $\alpha = 30^\circ, 60^\circ, 80^\circ$.

Furthermore, most parts of the proof suggest that the above argument might work with the Sobolev Space $W^{2,\infty}(K)$ (i.e “derivatives are Lipschitz continuous”).


Application II-2: Gravity-Driven Particle-Laden Flow

Optimal λ from leave-one-out cross validation:




Application II-2: Gravity-Driven Particle-Laden Flow

Quality of Approximation - Flux Function f :




Application II-2: Gravity-Driven Particle-Laden Flow

Quality of Approximation - Flux Function g :




Application II-2: Gravity-Driven Particle-Laden Flow

Quality of Approximation - Derivative of Flux Function f' :



Application II-2: Gravity-Driven Particle-Laden Flow

Quality of Approximation - Derivative of Flux Function g' :




Application II-2: Gravity-Driven Particle-Laden Flow

Riemann Initial Data:

$$(h, \phi_0)(0, x) = \begin{cases} (1, 0.4) & \text{for } x > 0, \\ (0.2, 0.4) & \text{for } x < 0. \end{cases}$$

Solution for $(h, \phi_0)(30, x)$:




Application II-2: Gravity-Driven Particle-Laden Flow

Riemann Initial Data:



$$(h, \phi_0)(0, x) = \begin{cases} (1, 0.485) & \text{for } x > 0, \\ (0.2, 0.485) & \text{for } x < 0. \end{cases}$$

Solution for $(h, \phi_0)(30, x)$:



Application II-2: Gravity-Driven Particle-Laden Flow

Violating Genuine Nonlinearity



Application II-2: Approximating Flux Functions

Computational Time for PDE Simulations,

$\Delta x = 0.001, \Delta t = 0.0005, t = 30, x \in [-0.1, 4],$

- Interpolation: 45s.
- Vectorized Polynomial Approximation: 984s.

Application II-2: Approximating Flux Functions

Computational Time for PDE Simulations,

$\Delta x = 0.001, \Delta t = 0.0005, t = 30, x \in [-0.1, 4],$

- Interpolation: 45s.
- Vectorized Polynomial Approximation: 984s.

Time to generate flux functions on a grid with $\Delta\phi_0 = 0.001$:

- 156s.

Application II-2: Approximating Flux Functions

Computational Time for PDE Simulations,

$\Delta x = 0.001, \Delta t = 0.0005, t = 30, x \in [-0.1, 4],$

- Interpolation: 45s.
- Vectorized Polynomial Approximation: 984s.

Time to generate flux functions on a grid with $\Delta\phi_0 = 0.001$:

- 156s.

Fix:

1. Generate sparse grid points.
2. Fit polynomials to f and g .
3. Pre-evaluate polynomials on a specified grid.

Application II-2: Approximating Flux Functions

Computational Time for PDE Simulations,

$\Delta x = 0.001, \Delta t = 0.0005, t = 30, x \in [-0.1, 4],$

- Interpolation: 45s.
- Vectorized Polynomial Approximation: 984s.

Time to generate flux functions on a grid with $\Delta\phi_0 = 0.001$:

- 156s.

Fix:

1. Generate sparse grid points.
2. Fit polynomials to f and g .
3. Pre-evaluate polynomials on a specified grid.
4. For any evaluation of f and g (especially in PDE simulations), perform numerical interpolation.

Application II-3: Comparing Models

Lubrication Assumption gives the **same system of conservation laws** for both diffusive flux and shear-induced migration models:

$$\begin{cases} h_t + \left(h^3 f \left(\frac{h\phi_0}{h} \right) \right)_x = 0, \\ (h\phi_0)_t + \left(h^3 g \left(\frac{h\phi_0}{h} \right) \right)_x = 0. \end{cases}$$

Application II-3: Comparing Models

Lubrication Assumption gives the **same system of conservation laws** for both diffusive flux and shear-induced migration models:

$$\begin{cases} h_t + \left(h^3 f \left(\frac{h\phi_0}{h} \right) \right)_x = 0, \\ (h\phi_0)_t + \left(h^3 g \left(\frac{h\phi_0}{h} \right) \right)_x = 0. \end{cases}$$

Different models yield different pairs of flux functions f and g .

Application II-3: Comparing Models

Lubrication Assumption gives the **same system of conservation laws** for both diffusive flux and shear-induced migration models:

$$\begin{cases} h_t + \left(h^3 f \left(\frac{h\phi_0}{h} \right) \right)_x = 0, \\ (h\phi_0)_t + \left(h^3 g \left(\frac{h\phi_0}{h} \right) \right)_x = 0. \end{cases}$$

Different models yield different pairs of flux functions f and g .

Observation: If the flux functions from different models are sufficiently close, solutions to the Riemann problems are sufficiently close!


Application II-3: Comparing Models

- Reference: A comparative study of dynamic models for gravity-driven particle-laden flows. (Lee W.P. et. al, 2025.)
- Authors: 2023 REU students, S.C. Burnett, L. Ding, A. L. Bertozzi.
- Accepted for publication in *Applied Mathematics Letters*, 2025.


Application II-3: Comparing Models

- Reference: A comparative study of dynamic models for gravity-driven particle-laden flows. (Lee W.P. et. al, 2025.)
- Authors: 2023 REU students, S.C. Burnett, L. Ding, A. L. Bertozzi.
- Accepted for publication in *Applied Mathematics Letters*, 2025.

$\alpha = 50^\circ$, Equilibrium Profile - I.




(a)




(b)

Application II-3: Comparing Models

$\alpha = 50^\circ$, Equilibrium Profile - II.




(c)




(d)

Application II-3: Comparing Models

$\alpha = 50^\circ$, PDE Simulations.



(b)



(c)

- **Main Result:** Under the usual assumptions and some additional mild assumptions, unique double-wave entropy solutions are preserved upon a sufficiently good approximation of flux functions.
- Understanding how each of the assumptions fails allows us to predict properties that are not expected to be preserved upon perturbation.

Generalizing the result to $n \times n$ systems.

- (Wong and Bertozzi, 2016.)
 $n = 3$: Bidensity/Bisize Particle Laden Flow
(Additional Parameter \rightarrow Additional Conservation Law.)
- General $n \times n$ using “more differential topology”.

Generalizing the result to $n \times n$ systems.

- (Wong and Bertozzi, 2016.)
 $n = 3$: Bidensity/Bisize Particle Laden Flow
(Additional Parameter \rightarrow Additional Conservation Law.)
- General $n \times n$ using “more differential topology”.

Other Variants - Regularity:

- Lower Regularity required for flux functions and their perturbations.
- Smooth except at finite points (corresponding to phase transitions).
- Perturbations to initial data (left and right states).

Allowing Linear Degenerate Waves:

- Example: $n = 3$, Compressible Euler Equations for gas dynamics.
- Expectation: Preserved under the class of “shocks, rarefactions, and contact discontinuities” for a class of perturbations.

Allowing Linear Degenerate Waves:

- Example: $n = 3$, Compressible Euler Equations for gas dynamics.
- Expectation: Preserved under the class of “shocks, rarefactions, and contact discontinuities” for a class of perturbations.

Violating Genuine Non-linearity:

- (Liu, 1973.)
Alternative to Lax's Entropy Condition \rightarrow Liu's Entropy Condition.
- Generalizing the above arguments for a different entropy condition.

Allowing Linear Degenerate Waves:

- Example: $n = 3$, Compressible Euler Equations for gas dynamics.
- Expectation: Preserved under the class of “shocks, rarefactions, and contact discontinuities” for a class of perturbations.

Violating Genuine Non-linearity:

- (Liu, 1973.)
Alternative to Lax’s Entropy Condition \rightarrow Liu’s Entropy Condition.
- Generalizing the above arguments for a different entropy condition.

Numerical Schemes Motivated by Transversality.



Thank you for your attention!



Particle-Laden Flow:



N. Murisic, J. Ho, V. Hu, P. Latterman, T. Koch, K. Lin, M. Mata, and A. L. Bertozzi.

Particle-laden viscous thin-film flows on an incline: Experiments compared with a theory based on shear-induced migration and particle settling.

Physica D: Nonlinear Phenomena 240, 20 (2011), 1661–1673.



N. Murisic, B. Pausader, D. Peschka, and A. L. Bertozzi.

Dynamics of particle settling and resuspension in viscous liquid films.

Journal of Fluid Mechanics 717 (2013), 203–31.



WP. Lee, J. Woo, L. Triplett, Y. Gu, S. C. Burnett, L. Ding, A. L. Bertozzi.

A comparative study of dynamic models for gravity-driven particle-laden flows.

Applied Mathematics Letters, 64, 109480 (2025).



J.T. Wong, A. L. Bertozzi

A conservation law model for bidensity suspensions on an incline.

Physica D: Nonlinear Phenomena 330 (2016), 47-57.



L. Wang, A. L. Bertozzi.

Shock solutions for high concentration particle-laden thin films.

SIAM J. Appl. Maths 74 (2014), 322–344.

References for Conservation Laws I:



H. Holden, L. Holden.

First-order nonlinear scalar hyperbolic conservation laws in one dimension.

Ideas and Methods in Mathematical Analysis, Stochastics, and Applications, 20 (1992), 480-510.



S. Bianchini, R. M. Colombo.

On the Stability of the Standard Riemann Semigroup

Proceedings of the American Mathematical Society 130 (2002), 1961–1973.



S. Schechter, D. Marchesin, B.J. Plohr.

Structurally stable Riemann solutions.

Journal of Differential Equations 126 (1996), 303-354.



A. Azevedo, C. Eschenazi, D. Marchesin, C. Palmeria.

Topological Resolution of Riemann Problems for Pairs of Conservation Laws

Quarterly of Applied Mathematics 68 (2010), 375-393.



C. Eschenazi, W. Lambert, M. Lopez-Flores, D. Marchesin, C. Palmeira.

Solving Riemann Problems with a Topological Tool (Extended version)

Journal of Differential Equations, 416 (2025), 2134–2174

References for Conservation Laws II:



D. Schaeffer.
A Regularity Theorem for Conservation Laws
Advances in Mathematics 11 (1973), 368-386.



L. Caravenna, L. V. Spinolo.
Schaeffer's Regularity Theorem for Scalar Conservation Laws Does Not Extend to Systems
Indiana University Mathematics Journal 66 (2017), 101-160.



T. Liu.
The Riemann Problem for General 2×2 Conservation Laws
Transactions of the American Mathematical Society 199 (1974), 89-112.



A. Bressan, G. Chen, S. Huang.
Generic singularities for 2D pressureless flows
Science China Mathematics 68 (2024) 559-576.

References for Conservation Laws III:



H. C. Kranzer, B. L. Keyfitz.

A strictly hyperbolic system of conservation laws admitting singular shocks
Nonlinear Evolution Equations That Change Type (1990), 107-125.



H. C. Kranzer, B. L. Keyfitz.

Existence and uniqueness of entropy solutions to the Riemann problem for hyperbolic systems of two nonlinear conservation laws
Journal of Differential Equations 27 (1978), 444-476.



B. Wendroff.

The Riemann Problem for Materials with Nonconvex Equations of State I:
Isentropic Flow
Journal of Mathematical Analysis and Applications 38 (1972), 454-466.



B. Wendroff.

The Riemann Problem for Materials with Nonconvex Equations of State II:
General Flow
Journal of Mathematical Analysis and Applications 38 (1972), 640-658.

Differential Topology:



V. Guillemin, A. Pollack.
Differential Topology
Prentice-Hall, 1974.



M. W. Hirsch.
Differential Topology
Springer New York, NY, 1972.



C. Greenblatt.
An Introduction to Transversality.
https://schapos.people.uic.edu/MATH549_Fall2015_files/Survey%20Charlotte.pdf,
2015. Accessed 25 March 2024.



J. M. Lee.
Introduction to Smooth Manifolds.
Springer New York 2nd Edition, 2012.



S. Ichiki.
Characterization of Generic Transversality
Bulletin of the London Mathematical Society 51 (2019), 978-988.

Other Textbooks/Monographs:



P. D. Lax.
Hyperbolic Systems of Conservation Laws II
Communications on Pure and Applied Mathematics 10 (1957), 537-566.



R. J. LeVeque.
Numerical Methods for Conservation Laws.
Birkhäuser Basel, 2nd ed, 1992.



H. Holden , N. H. Risebro
Front Tracking for Hyperbolic Conservation Laws.
Applied Mathematical Sciences 152, 2nd ed, 2015.



J. K. Hunter, B. Nachtergaele.
Applied Analysis.
World Scientific Publishing Co, 2001.